

*

دانشکده مهندسی شیمی-پردیس دانشکده‌های فنی - دانشگاه تهران

Huron-Vidal (PR) (SRK)
 Huron-Vidal) C₂
 Chrastil Mendez

Chrastil Mendez

Charstil, Aguilera, Gordillo, Mendez

SRK, PR

.[]

(SFE)

: [] SFE

SFE

()

[]

[]

(%AARD)

$$AARD(\%) = \frac{1}{n} \left| \sum \frac{y_{2,exp} - y_{2,calv}}{y_{2,exp}} \right| \times 100 \quad ()$$

n

[]

[]

[]

Solute	M [g/mol]	P _c [MPa]	T _c [K]	ω	A	B [K]	V ^{sol} *10 ³ [m ³ /mol]
Phenanthrene	178.24	3.17	882.55	0.3299	14.631	4873.4	0.1512
Benzoic Acid	122.124	4.50	752.00	0.620	14.408	4618.1	0.0965
Hexachloroethane	236.74	3.45	714.60	0.1630	10.6322	2600.94	0.1132
Triphenylmethane	244.34	2.24	863.00	0.5760	14.7858	5228.0	0.2409
Naphthalene	128.174	4.05	748.15	0.3020	13.583	3733.9	0.1103
2,3-dimethylnaphthalene	156.23	3.22	785.00	0.4240	14.0646	4302.5	0.1547
2,6-dimethylnaphthalene	156.23	3.22	777.00	0.4201	14.4286	4419.5	0.1547
Fluorene	166.23	2.99	821.00	0.4070	14.2046	4561.8	0.1393
Anthracene	178.234	3.12	869.30	0.3531	12.147	4397.6	0.1426
Pyrene	202.25	2.61	936	0.509			0.6299
Ascorbyl Palmitate	414.50	1.156	870.81	1.85	-	-	0.3405
BHA	180.2	2.883	798.78	0.63	-	-	0.1688
Dodecyl Gallate	338.45	1.846	905.90	1.20	-	-	0.2679
Propyl Gallate	170.12	4.772	862.87	0.86	-	-	0.155

Mendez $T \ln(yP) = A' + B' \rho + C'T$ A', B', C' \vdots $()$ $B' ()$ k \vdots $k = n_1 T + m_1$ $B' = n_2 T + m_2$ $()$ \vdots $\ln S_2 = (n_1 T + m_1) \ln \rho + \frac{\alpha}{T} + \beta$ $()$ \vdots $T \ln(y_2 P) = A' + (n_2 T + m_2) \rho + C'T$ \vdots $Eviews 3.1$	Chrastil \vdots B $k - A$ \vdots $\ln S_2 = k \ln \rho + \frac{\alpha}{T} + \beta$ β $\Delta H) \quad \frac{\Delta H}{R}$ \vdots $: []$ $y - S_2$ $S_2 = \frac{\rho M_{wsolute} y_2}{M_{wsolvent} (1 - y_2)}$ \vdots $\boxed{\text{Del valle and Aguilera}}$ \vdots $\ln S_2 = k \ln \rho + \frac{\alpha_1}{T} + \frac{\alpha_{11}}{T^2} + \beta$ $k, \alpha_1, \alpha_{11}, \beta$ \vdots $\boxed{\text{Del valle and Aguilera}} \quad \boxed{\text{Chrastil}}$ \vdots $()$ \vdots $\ln(y_2) = D_0 + D_1 P + D_2 P^2 + D_3 PT + D_4 T + D_5 T^2$ $D_0, D_1, D_2, D_3, D_4, D_5$ \vdots $\boxed{\text{Mendez-Santiago and Teja}}$
--	--

$a_c = \eta_2 \frac{R^2 T_c^2}{P_c}$	()	Chrastil
$\alpha(T_r) = [1 + m(1 - \sqrt{T_r})]^2$	()	Chrastil
$m_1 = \lambda_1 + \lambda_2 \omega + \lambda_3 \omega^2$	()	
$\eta_1, \eta_2, \lambda_1, \lambda_2, \lambda_3$	Aguilera	

(GCM) () ()


$$y_2 = \frac{f_2^{os}}{\phi_2^v P} = \frac{P_2^{Sub}}{P} \frac{\exp(\frac{\nu_2^s(P - P_2^{Sub})}{RT})}{\phi_2^v} \quad ()$$

Huron and Vidal


C₂ PR SRK

$$\begin{aligned} & P = \frac{RT}{v-b} - \frac{a}{v(v+b)} \quad () \\ & P = \frac{RT}{v-b} - \frac{a}{v(v+b)+b(v-b)} \quad () \\ & B_m = \sum y_i B_i = y_1 B_1 + y_2 B_2 \quad () \quad b, a \\ & \frac{A_m}{B_m} = \sum y_i \left(\frac{A_i}{B_i} - \frac{\ln \gamma_i^\infty}{\ln 2} \right) = \sum y_i C_i \quad () \\ & C_2 = \frac{A_2}{B_2} - \frac{\ln \gamma_2^\infty}{\ln 2} \quad () \quad b = \eta_1 R \frac{T_c}{P_c} \quad () \\ & a(T) = a_c \alpha(T_r) \quad () \end{aligned}$$


	Eq. (2)	Eq. (4)	Eq. (5)	Eq. (6)	Eq. (9)	Eq. (10)
Anthracene- CO ₂	18.04	14.02	37.59	29.30	17.83	28.04
Anthracene-Ethane	9.94	9.47	7.09	13.75	9.30	13.11
Fluorene- CO ₂	21.72	17.30	73.85	32.93	21.70	32.52
Fluorene-Ethylene	8.10	6.67	49.97	35.75	6.91	21.87
Naphthalene-Ethane	13.93	14.27	83.96	16.19	11.40	15.43
Triphenylmethane- CO ₂	9.65	7.32	21.49	10.87	8.90	10.94
Pyrene- CO ₂	23.43	23.38	59.24	44.54	20.28	34.78
Pyrene-Ethylene	7.98	-	-	34.99	7.93	22.12
Phenanthrene-CO ₂	5.86	5.74	10.12	9.44	4.28	5.15
Phenanthrene-C ₂ H ₄	11.15	9.04	8.92	10.22	10.63	7.34
Phenanthrene-Ethylene	2.55	1.34	0.00	1.58	1.39	1.27
Dodecyl Gallate- CO ₂	5.58	-	-	6.28	1.50	2.61
Propyl Gallate- CO ₂	3.60	-	-	4.85	2.29	2.28
Ascorbyl Palmitate- CO ₂	5.07	-	-	3.77	-	3.38
Butyl Hydroxyl Anisole- CO ₂	8.30	-	0.00	10.06	3.78	4.74
2,6-Dimethylnaphthalene-C ₂ H ₄	10.03	9.37	18.35	12.80	10.39	6.40
2,6-Dimethylnaphthalene-CO ₂	9.38	9.36	23.16	15.54	8.89	9.01
2,3-Dimethylnaphthalene-CO ₂	13.19	13.00	21.32	18.00	11.65	15.08
Benzoic acid-CO ₂	9.81	9.67	11.74	16.16	9.22	9.54
Hexachloroethane-CO ₂	15.30	15.29	16.99	15.63	14.53	10.26
Benzoic acid-C ₂ H ₄	8.88	9.01	7.93	12.42	8.86	11.69
%Average	12.07	11.71	34.89	19.89	11.17	15.70




PR-EOS




SRK-EOS




PR-EOS




SRK-EOS




SRK-EOS



PR-EOS



PR-EOS



SRK-EOS

PR-EOS () :					
	a_1	a_2	b_1	b_2	P_{ref}
Anthracene- CO ₂	-11.71061	0.15181	-48.41242	0.15445	90.6
Anthracene-Ethane	63.95572	-0.06400	13.74482	0.04552	104.3
Fluorene- CO ₂	31.57406	0.01874	-22.25118	0.07381	69.9
Fluorene-Ethylene	119.24868	-0.21770	11.15511	-0.02282	69.9
Naphthalene-Ethane	26.87023	0.03461	-38.78105	0.13394	35.4
Pyrene- CO ₂	53.40834	0.00055	-14.19848	0.05861	104.3
Pyrene-ethylene	126.54046	-0.19651	12.68261	-0.02073	104.8
Phenanthrene- CO ₂	0.22600	-67.38105	0.11201	11.77612	80.9
Phenanthrene-C ₂ H ₄	77.64750	-0.06460	-32.58310	0.12384	120
Phenanthrene-Ethlene	106.85941	-0.24870	50.96796	-0.17492	276.8
Triphenylmethane-Ethane	56.221648	-0.00221	-72.679333	0.24111	69.9
Ascorbyl Palmitate- CO ₂	44.22489	-0.10575	271.25330	-0.91581	130
BHA- CO ₂	-46.75261	0.28427	-198.29651	0.62973	150
Dodecyl Gallate- CO ₂	-363.61649	1.33476	-293.29176	0.95057	150
Propyl Gallate- CO ₂	-221.55230	0.82812	-224.46840	0.71513	150
2,3-Dimethylnaphthalene- CO ₂	5.58520	0.11695	-22.60553	0.07780	99
2,6-Dimethylnaphthalene- C ₂ H ₄	120.13071	-0.22341	17.76860	-0.03965	80
2,6-Dimethylnaphthalene- CO ₂	-12.20245	0.17150	-89.03309	0.297778	97
Benzoic acid- C ₂ H ₄	98.70964	-0.14387	-4.21413	0.00493	120
Benzoic acid- CO ₂	47.35324	0.01744	-35.37715	0.13352	120
Hexachloroethane- CO ₂	-5.43943	0.10441	-46.73699	0.15133	99

SRK-EOS () :					
	a_1	a_2	b_1	b_2	P_{ref}
Anthracene- CO ₂	22.36295	-0.00874	-10.98372	0.03450	90.6
Anthracene-Ethane	40.01947	-0.05203	-3.94254	0.01390	104.3
Fluorene- CO ₂	35.06430	-0.04513	-5.16126	0.01858	69.9
Fluorene-Ethylene	64.41210	-0.11899	6.78220	-0.01549	69.9
Naphthalene-Ethane	13.89840	0.02383	-19.27564	0.06826	35.4
Pyrene- CO ₂	53.03783	0.00055	-14.19848	0.05861	104.3
Pyrene-ethylene	64.38652	-0.09852	2.62839	0.00007	104.8
Phenanthrene- C ₂ H ₄	48.06919	-0.05757	-6.56959	0.02949	120
Phenanthrene-CO ₂	0.05001	-13.31905	-0.02305	33.15711	80.9
Phenanthrene-Ethlene	58.23523	-0.13545	28.75895	-0.09792	276.8
Triphenylmethane-Ethane	48.62361	-0.05926	-20.99134	0.07141	69.9
Ascorbyl Palmitate- CO ₂	25.88340	-0.06179	138.06860	-0.46581	130
BHA- CO ₂	-18.73836	0.13650	-72.01950	0.22990	150
Dodecyl Gallate- CO ₂	-129.90112	0.51080	-80.72521	0.26513	150
Propyl Gallate- CO ₂	-79.57368	0.31978	-73.42003	0.233622	150
2,3-Dimethylnaphthalene- CO ₂	30.73873	-0.02595	-22.60553	0.077795	99
2,6-Dimethylnaphthalene- C ₂ H ₄	60.94417	-0.11113	8.87989	-0.021756	80
2,6-Dimethylnaphthalene- CO ₂	18.41990	0.01194	-25.81840	0.08807	97
Benzoic acid- C ₂ H ₄	55.57447	-0.08914	6.67402	-0.01351	120
Benzoic acid- CO ₂	52.18188	-0.07855	9.16441	-0.02103	120
Hexachloroethane- CO ₂	11.77513	0.01132	-7.38065	0.02559	99

Chrastil

(% /)

Gordillo

(% /)

Chrastil

$n_1 T \ln \rho$

Aguilera

(C₂)

C₂

C₂

C₂

:T		
:Tc		$:f_2^{0s}$
:Tr		:Mw
$:V_2^s$:N
$:y_2$:P
i	$\hat{\phi}_i$:Pc
	$:P_2^{Sub}$:Pr
	$:R$	
	$:S_2$	

()		:
	SRK-EOS	PR-EOS
Anthracene- CO ₂	17.22	15.73
Anthracene-Ethane	7.78	8.18
Fluorene- CO ₂	10.02	11.59
Fluorene-Ethylene	21.07	30.16
Naphthalene-Ethane	30.34	42.44
Pyrene- CO ₂	16.41	26.07
Pyrene-ethylene	12.93	15.92
Phenanthrene- C ₂ H ₄	15.74	20.25
Phenantrene- CO ₂	9.87	9.30
Phenantrene-Ethylene	3.02	2.94
Triphenylmethane- CO ₂	34.38	43.33
Ascorbyl Palmitate- CO ₂	6.32	6.41
BHA- CO ₂	16.50	29.52
Dodecyl Gallate- CO ₂	7.12	10.59
Propyl Gallate- CO ₂	3.98	6.00
2,3-Dimethylnaphthalene- CO ₂	8.86	22.62
2,6-Dimethylnaphthalene- C ₂ H ₄	18.01	26.66
2,6-Dimethylnaphthalene- CO ₂	10.79	22.13
Benzoic acid- C ₂ H ₄	3.35	4.29
Benzoic acid- CO ₂	5.93	13.53
Hexachloroethane- CO ₂	6.58	23.41
%AARE	14.00	19.85

%AARD :				
	Mendez	Chrastil	SRK-EOS	PR-EOS
Coimbra	6.3	7.9	28.14	24.65
	6.2	6.9	13.70	18.15

Equation of state		Chrastil model	
PR-EOS	SRK-EOS	Modified form	Main form
%AARD	17.01	8.97	5.99
			6.40

- 1- Lee, L.S., Fu, J.H., and Hsu, H.L. (2000). "Solubility of solid 1,4-dimethoxybenzene in supercritical carbon dioxide." *J. Chem. Eng. Data*, Vol. 45, PP. 358-361.
- 2- Jouyban, A., Chan, H.K., and Foster, N.R. (2002). "Mathematical representation of solute solubility in supercritical carbon dioxide using empirical expressions." *J. of Supercritical Fluids*, Vol. 24, PP. 19-35.
- 3- Khimechea, K., Alessi, P., Kikic, I., and Dahmani, A. (2007). "Solubility of diamines in supercritical carbon dioxide experimental determination and correlation." *J. of Supercritical Fluids*, Vol. 41, PP. 10-19.

-
- 4- Saldana, M., Tomberli, B., Selma E., Goldman, G.S., Gray, C.G., and Temelli, F. (2007). "Determination of vapor pressure and solubility correlation of phenolic compounds in supercritical CO₂." *J. of Supercritical Fluids*, Vol. 40, PP. 7–19.
- 5- Abolghasemi, H., Orouj, R., Tabasi, M., (2006) "The study of effective parameters on metals extraction (Cu, Co, Hg, and U) with supercritical fluid." *Iranian Association of Chemical Engineering*, Vol.23, PP.22-30.
- 6- Johnston, K.P., Zlger, D.H., and Eckert, C.A. (1982). "Solubilities of hydrocarbon solids in supercritical fluids. The augmented van der waals treatment." *Ind. En\$. Chem. Fundam*, Vol. 21, PP. 191-197.
- 7- Murga, R., Sanaz, M.T., Beltran, S., and Cabezas, J.L. (2003). "Solubility of three hydroxycinnamic acids in supercritical carbon dioxide." *J. of Supercritical Fluids*, Vol. 27, PP. 239-245.
- 8- Ozel, M.Z., Bartle, K.D., Clifford, A.A., and Burford, M.D. (2000) "Extraction, solubility and stability of metal complexes using stainless steel supercritical fluid extraction system." *Analytica Chimica Acta*, Vol. 417, PP. 172-184.
- 9- Cross, W., Akgerman, J.A., and Erkey, C. (1996). "Determination of metal-chelate complex solubilities in supercritical carbon dioxide." *Ind. Eng. Chem. Res*, Vol. 35, PP. 1765-1770.
- 10- Li, H., and Yang, S.X. (2003). "Modelling of supercritical fluid extraction by hybrid Peng-Robinson equation of state and genetic algorithms." *Biosystems Engineering*, Vol. 86, PP. 17-25.
- 11- Orouj, R., (2007) "Investigation on extraction of metal elements from solid matrix with supercritical fluid." *Thesis for MSc Degree*, Tehran University.
- 12- Jin, J., Zhong, C., Zhang, Z., and Li, Y. (2004). "Solubilities of benzoic acid in supercritical CO₂ with mixed cosolvent." *Fluid Phase Equilibria*, Vol. 226, PP. 9–13.
- 13- Cortesi, A., Kikic, I., Alessi, P., Turtoi, G., and Garnier, S. (1999). "Effect of chemical structure on the solubility of antioxidants in supercritical carbon dioxide: experimental data and correlation." *J. of Supercritical Fluids*, Vol. 14, PP. 139–144.
- 14- Kurnlk, R.T., Holla, S.J., and Reid R.C. (1981). "Solubility of solids in supercritical carbon dioxide and ethylene." *J.Chem. Eng. Data*, Vol. 26, PP. 47-51.
- 15- Chrastil, F.J. (1982). "Solubility of solids and liquids in supercritical gases." *phys. Chem*, Vol. 86, PP. 3016-3021.
- 16- del Valle, J.M., and Aguilera, J.M. (1988). "An improved equation for predicting the solubility of vegetable oils in supercritical CO₂." *Industrial and Engineering Chemistry Research*, Vol. 27, PP. 1551–1553.
- 17- Gordillo, M. D., Blanco, M. A., Molero, A., and Martinez de la Ossa, E. (1999). "Solubility of the antibiotic Penicillin G in supercritical carbon dioxide." *J. of Supercritical Fluids*, Vol. 15, PP. 183–190.
- 18- Mendez-Santiago, J., Teja, A.S. (2000). *Ind. Eng. Chem. Res.*, Vol. 39, PP. 4767–4771.
- 19- Orouj, R., Abolghasemi, H., Mahdavian, M., and Vatani, Z. (2007) "Prediction solubility of spiroindolinonaphthoxazine photochromic dye in supercritical carbon dioxide by using equation of states." *Iranian Journal of Chemistry and Chemical Engineering*, Vol. 26.
- 20- Coimbra, P., Gil, M.H., Duarte, C.M.M., Heron, B.M., and de Sousa, H.C. (2005). "Solubility of a Spiroindolinonaphthoxazine photo chromic dye in Supercritical carbon dioxide: Experimental determination and correlation." *Fluid Phase Equilibria*, Vol. 238, PP. 120-128.
-

-
- 21- Orouj, R., Abolghasemi, H., Tabasi, M., and Mahdavian, M. (2007). "Predicting solubility of spiroindolinonaphthoxazine photochromic dye in supercritical carbon dioxide by using equations of state." *5th International symposium in chemical engineering and high pressure processes*, Valladolid, Spain.
- 22- Orouj, R., Abolghasemi, H., Tabasi, M., and Mahdavian, M. (2007). "Predicting solubility of tributyl phosphate in supercritical carbon dioxide." *European Congress of Chemical Engineering (ECCE-6)* Copenhagen.

- 1- Super Critical Fluid Extraction
2- Absolute Average Relative Deviation
3- Group Contribution Method